

HANDLE MOVES IN CONTACT SURGERY DIAGRAMS

FAN DING AND HANSJÖRG GEIGES

ABSTRACT. We describe various handle moves in contact surgery diagrams, notably contact analogues of the Kirby moves. As an application of these handle moves, we discuss the respective classifications of long and loose Legendrian knots.

1. INTRODUCTION

In [3] and [4] we introduced the notion of contact Dehn surgery, especially contact (± 1) -surgery, see also [16, Section 6.4]. The resulting contact surgery diagrams were studied more carefully in [6] and since then have led — in particular when combined with Heegaard Floer theory — to a wealth of applications, see e.g. [17], [26], [30]. In the present paper we discuss a variety of handle moves in such contact surgery diagrams: first and second Kirby move, handle cancellation, and handle moves involving 1-handles translated into contact $(+1)$ -surgeries. We hope that this will add to the usefulness of contact surgery diagrams.

The second Kirby move (of which we shall present three ‘incarnations’) is fundamental to most of the results in this paper. A further important ingredient is the Legendrian isotopy between the push-off and the meridian of a (-1) -surgery curve. These two results are proved using convex surface theory.

Two applications of such handle moves (or the methods used in their proof) will be discussed: the relation between the classification of long Legendrian knots in $\mathbb{R}^3$ and their closures in S^3 , and the classification of loose Legendrian knots, i.e. knots with overtwisted complements. The first is in response to a question by D. Fuchs and S. Tabachnikov, the second extends a result of K. Dymara.

We assume familiarity with the fundamental notions of contact topology on the level of [16]. Our contact structures are understood to be coorientable, i.e. they can be written as $\xi = \ker \alpha$ with $\alpha \wedge d\alpha \neq 0$. With ξ_{st} we denote the standard contact structure $\ker(dz + x dy)$ on $\mathbb{R}^3$ as well as the standard contact structure on S^3 . Legendrian knots are always illustrated via their front projections in the yz -plane.

2. LEGENDRIAN BAND-SUM AND THE SECOND KIRBY MOVE

In this section we formulate the contact analogue of a 2-handle slide, or what is also called a Kirby move of the second kind.

By a *Legendrian band* we mean a band in $(\mathbb{R}^3, \xi_{\text{st}})$ whose boundary curves consist of an arbitrary Legendrian curve and its Legendrian push-off in the z -direction. Given two Legendrian knots L_1 and L_2 in $(\mathbb{R}^3, \xi_{\text{st}})$, we can form their *Legendrian band-sum* $L_1 \#_{\text{Lb}} L_2$. This means that we construct the connected sum with the help of a Legendrian band. Here any band disjoint from the rest of the link may be used. If we think of L_1 and L_2 as surgery curves, then — topologically — the

replacement of L_1, L_2 by $L_1 \#_{\text{Lb}} L_2, L_2$ corresponds to a handle slide of the 2-handle corresponding to L_1 over that corresponding to L_2 . If L_1 and L_2 are oriented, one can sensibly speak of *handle addition* or *handle subtraction*; see Figure 1, where we write $\overline{L}_2$ for L_2 with reversed orientation (and the knots are drawn as trivial knots, but the picture is meant to represent arbitrary knots, which may also be linked).




FIGURE 1. 2-handle addition and subtraction.

Remark. There is an alternative way for forming the connected sum of two Legendrian knots in the front projection picture, see [14, Figure 2]. This could also be used here. We shall return to this point in Section 8.

After a topological handle slide, one has to compute the new surgery framing of $L_1 \# L_2$ from the original surgery framings and the linking number of L_1 and L_2 (in the case of handles attached to the 4-ball D^4), see [20, p. 142]. Here it is obviously necessary to keep track of the orientation of L_2 , and to make the distinction between handle addition and subtraction. For handles attached to the boundary of an arbitrary 4-manifold, one can compute the new framing with the help of the double-strand notation for framings.

In the contact geometric setting it turns out that, relative to the contact framing, we do not have to compute new surgery framings at all; in particular, we may allow rational surgery framings for L_1 , although only integer framings correspond to actual topological handle slides. (Beware, though, that contact surgery with rational coefficient different from $1/k$ with $k \in \mathbb{Z}$ is not uniquely defined, so one can only say that each particular contact r -surgery along L_1 corresponds to an r -surgery along a Legendrian isotopic copy of L_1 .) Notice, moreover, that one can make sense of a Legendrian band in an arbitrary contact manifold, and we shall formulate the following proposition accordingly.

Proposition 1 (The second Kirby move). *Let L_1, L_2 be two Legendrian knots in a contact 3-manifold (M, ξ) . Let $L_2^\pm$ be the Legendrian knot obtained as a push-off of L_2 with one additional positive or negative twist, see Figure 2. Then, in the manifold obtained from (M, ξ) by contact (± 1) -surgery along L_2 , the knot L_1 is Legendrian isotopic to $L_1 \#_{\text{Lb}} L_2^\pm$.*

Proof. By a *Legendrian unknot* L_0 we shall mean a Legendrian knot that can be represented in the front projection of a suitable Darboux chart by a front with two




FIGURE 2. Adding a positive or negative twist.

cusps and no crossings. The Legendrian band-sum with such a Legendrian unknot is trivial, provided the band is disjoint from a disc bounded by L_0 : the Legendrian isotopy between $L \#_{\text{Lb}} L_0$ and the original Legendrian knot L is given, in Darboux charts, by a sequence of first and second Legendrian Reidemeister moves (see [32] or [12] for these moves). An instance of this phenomenon can be read off from Figure 1.

Therefore, in order to prove the proposition, it suffices to show that in the manifold obtained by contact (± 1) -surgery along L_2 , the knot $L_2^\pm$ is a Legendrian unknot as described.

We are going to write T for any 2-torus that forms the boundary of a tubular neighbourhood νL_2 of L_2 . For the computation of framings we are about to perform, the specific choice of T will be irrelevant. A Legendrian push-off L'_2 of L_2 can be realised as one of two Legendrian divides on a convex torus T . Along this Legendrian divide, the contact planes are tangent to T , so the twisting $\mathfrak{t}_T(L'_2)$ of the contact planes along L'_2 relative to the framing induced by T equals 0.

The knots L'_2 and $L_2^\pm$ are topologically isotopic, so we can compare their framings induced by ξ and T , respectively. As usual, right-handed twists are counted positively. Then the ξ -framing of $L_2^\pm$ relative to the ξ -framing of L'_2 equals $\pm 1 - 1$; the T -framing of $L_2^\pm$ relative to the T -framing of L'_2 equals ± 1 . It follows that

$$\mathfrak{t}_T(L_2^\pm) = \mathfrak{t}_T(L'_2) - 1 = -1.$$

In the surgered manifold, $L_2^\pm$ becomes a meridian of the solid torus we glue in to replace the solid torus νL_2 . (In particular, $L_2^\pm$ is homologically trivial in the surgered manifold, and we can speak of its Thurston-Bennequin invariant $\mathfrak{tb}(L_2^\pm)$ there.) So here the surface framing of $L_2^\pm$ given by T coincides with that given by a meridional disc D with boundary $L_2^\pm$. This implies that

$$\mathfrak{tb}(L_2^\pm) = \mathfrak{t}_D(L_2^\pm) = \mathfrak{t}_T(L_2^\pm) = -1$$

in the surgered manifold. Observe that D is disjoint from the Legendrian band used to connect L_1 and $L_2^\pm$ in M .

Moreover, $L_2^\pm$ can be chosen arbitrarily close to L_2 . This guarantees that after the surgery it is a topologically trivial knot inside a solid torus with tight contact structure. By the theorem of Eliashberg and Fraser [10, 11], $L_2^\pm$ is a Legendrian unknot in the surgered manifold. $\square$

Remark. Variants of the second Kirby move will be discussed in Section 8 in the context of the so-called ‘move 6’ for Legendrian knots, one of the Reidemeister moves in the presence of 1-handles.

3. PUSH-OFF AND MERIDIAN OF (-1) -SURGERY CURVE

The aim of the present section is to prove the following proposition, which will be instrumental in showing how to replace 1-handles by contact $(+1)$ -surgeries.

Proposition 2. *In the contact manifold obtained from any other contact 3-manifold (M, ξ) by contact (-1) -surgery along a Legendrian knot L , the Legendrian push-off λ of L and a standard Legendrian meridian μ (as shown in Figure 3) are Legendrian isotopic.*




FIGURE 3. λ is Legendrian isotopic to μ in the surgered manifold.

Proof. (i) We may assume that λ and μ intersect transversely in a single point. Choose a tubular neighbourhood νL of L such that $\mu, \lambda \subset \partial(\nu L)$. We claim that there is a C^0 -small isotopy of $\partial(\nu L)$ rel μ, λ , turning it into a convex surface, with characteristic foliation as depicted in Figure 4; the dashed lines indicate the dividing curves.




FIGURE 4. The characteristic foliation on $\partial(\nu L)$.

(ii) Assuming this claim for the time being, the proof concludes as follows. Write $\mu_0 := * \times \partial D^2$, $* \in S^1$, and $\lambda_0 := S^1 \times *$, $* \in \partial D^2$, for meridian and longitude, respectively, of the solid torus $S^1 \times D^2$ that we glue in replacing νL when we perform surgery along L . The attaching map for a contact (-1) -surgery can be described by

$$\mu_0 \mapsto \mu - \lambda, \quad \lambda_0 \mapsto \mu.$$

In other words, we may assume this gluing map to send λ_0 to a Legendrian ruling curve on $\partial(\nu L)$, and $\lambda_0 - \mu_0$ to a Legendrian divide. It follows that the contact structure on the glued in $S^1 \times D^2$, giving us the extension of the contact structure ξ on $M \setminus \text{Int}(\nu L)$ to that on the surgered manifold, is given — up to isotopy rel boundary — by the standard model

$$\cos \theta dx - \sin \theta dy = 0.$$

In this model, the two Legendrian divides $\lambda_0 - \mu_0$ as well as the Legendrian ruling curves λ_0 are Legendrian isotopic to the spine $S^1 \times \{0\}$.

(iii) It remains to prove the claim from (i). Write $\Sigma := \partial(\nu L)$, and identify this 2-torus with

$$\mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} \times \{0\} \subset \mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z} \times \mathbb{R},$$

with μ, λ corresponding to the first and second factor $\mathbb{R}/\mathbb{Z}$, respectively. Write x, y, z for the coordinates corresponding to these three factors. The twisting $\mathfrak{t}_\Sigma(\lambda)$ of the contact planes along λ relative to the framing induced by Σ equals

$$\mathfrak{t}_\Sigma(\lambda) = 0,$$

since λ may be realised as one of two Legendrian divides on a standard convex torus Σ with two dividing curves; along such a Legendrian divide the contact planes coincide with the tangent planes to Σ . Furthermore, with D denoting a meridional disc with $\partial D = \mu$, we have

$$\mathfrak{t}_\Sigma(\mu) = \mathfrak{t}_D(\mu) = \mathfrak{tb}(\mu) = -1.$$

It follows that we may perturb Σ rel μ, λ in such a way that the contact structure near the one-point union of μ and λ looks like

$$\sin(2\pi x) dy + \cos(2\pi x) dz = 0.$$

So the characteristic foliation looks as in Figure 5 (a), with singular points drawn in bold face.




FIGURE 5. The characteristic foliation on $\Sigma = \partial(\nu L)$ after the first and second perturbation, respectively.

Now perturb Σ relative to a small neighbourhood of μ and λ such that ξ looks like

$$(1) \quad a(x, y) dx + \sin(2\pi x) dy + \cos(2\pi x) dz = 0,$$

with a smooth function $a(x, y)$ satisfying the following conditions (for some small $\varepsilon > 0$):

- $a(x, y) = 0$ for $y \leq \varepsilon$ or $y \geq 1 - \varepsilon$ or $x \leq \varepsilon$ or $x \geq 1 - \varepsilon$,
- $a(x, y) < 0$ for x near $1/2$ and $y \in (\varepsilon, 2\varepsilon)$,
- $a(x, y) \leq 0$ for $y \in [0, 2\varepsilon]$,
- analogous conditions near $y = 1$ with reversed sign of $a(x, y)$,
- $\frac{\partial a}{\partial y}$ sufficiently small for (1) to define a contact structure.

(The conversion from such a perturbation of ξ to a perturbation of Σ with fixed ξ is a standard argument using Gray stability, see [16, Prop. 4.6.11].)

Then the characteristic foliation on Σ is given by the vector field

$$X := \sin(2\pi x) \partial_x - a(x, y) \partial_y,$$

as indicated in Figure 5 (b). Now perturb Σ in the interior (i.e. away from μ and λ) to make the characteristic foliation of Morse-Smale type there. This is possible by the density theorem of Kupka and Smale, see [29] or [25]. In fact, in the present situation the vector field X defining the characteristic foliation near μ and λ is transverse to the boundary of a disc-shaped region in the interior, and one may appeal directly to somewhat more elementary results of Peixoto [28] about Morse-Smale vector fields on surfaces with boundary, with the vector field being transverse to the boundary.

The resulting surface Σ will be convex, cf. [16, Section 4.8]. With Γ denoting its dividing set, a result of Kanda [24] allows us to compute the geometric number of intersection points of μ and λ , respectively, with Γ as follows:

$$-\frac{1}{2}\#(\mu \cap \Gamma) = \mathbf{t}_\Sigma(\mu) = -1, \quad -\frac{1}{2}\#(\lambda \cap \Gamma) = \mathbf{t}_\Sigma(\lambda) = 0.$$

So the dividing set must consist of two curves parallel to λ . By a final perturbation of Σ (away from μ and λ) we may achieve the characteristic foliation depicted in Figure 4, since that foliation is divided by Γ . $\square$

4. TOPOLOGICALLY TRIVIAL LONG LEGENDRIAN KNOTS

A *long Legendrian knot* is a Legendrian embedding $\gamma: \mathbb{R} \rightarrow (\mathbb{R}^3, \xi_{\text{st}})$ with $\gamma(t) = (0, t, 0)$ for $|t|$ large. By an isotopy of long knots we mean an isotopy that is stationary outside a compact subset of $\mathbb{R}$. A long knot is *topologically trivial* if it is isotopic to the embedding $t \mapsto (0, t, 0)$. We write $K = \gamma(S^1)$ for the unparametrised knot; we also use that notation if the particular parametrisation (except its orientation) is irrelevant.

The classical invariants $\mathbf{tb}$ (Thurston-Bennequin invariant) and $\mathbf{rot}$ (rotation number) of a long Legendrian knot are computed from its front projection in the same way as the invariants of an honest (‘short’) Legendrian knot. Any long Legendrian knot K can be completed to a short Legendrian knot $\hat{K}$ as shown in Figure 6. Then $\mathbf{tb}(\hat{K}) = \mathbf{tb}(K) - 1$ and $\mathbf{rot}(\hat{K}) = \mathbf{rot}(K)$.




FIGURE 6. A long knot K and its completion $\hat{K}$.

Alternatively, one can obtain the completion $\hat{K}$ by regarding K as a Legendrian knot in (S^3, ξ_{st}) passing through the point at infinity; by a Legendrian isotopy one can then move $\hat{K}$ into $\mathbb{R}^3 \subset S^3$. This is feasible because the complement of a point in (S^3, ξ_{st}) is contactomorphic to $(\mathbb{R}^3, \xi_{\text{st}})$, see [16, Prop. 2.1.8].

In [15, Section 2.5] the following result is mentioned in passing.

Proposition 3. *Topologically trivial long Legendrian knots are classified, up to Legendrian isotopy, by the two classical invariants.*

Proof. A very short proof of the corresponding result for topologically trivial Legendrian knots in (S^3, ξ_{st}) , originally due to Eliashberg and Fraser [10, 11], can be found in [13, Section 3.5]. The argument given there for showing that any topologically trivial Legendrian knot with $\text{tb} < -1$ destabilises carries over to long knots with $\text{tb} < 0$ without change. Only the proof that there is a unique topologically trivial long Legendrian knot with $\text{tb} = 0$ (the maximal possible value of tb in that case) requires a minor modification.




FIGURE 7. The characteristic foliation on the disc bounded by the trivial Legendrian knot.

Thus, let K be a topologically trivial *long* Legendrian knot in $(\mathbb{R}^3, \xi_{\text{st}})$ with $\text{tb}(K) = 0$. We regard its completion $\hat{K}$ as a Legendrian knot in (S^3, ξ_{st}) with $\text{tb}(\hat{K}) = -1$. Let D be a disc bounded by $\hat{K}$, which we may assume to be convex. Then the dividing set Γ_D consists of a single arc, for otherwise $\hat{K}$ could be destabilised, contradicting the maximality of tb . After a C^0 -small perturbation of D , the characteristic foliation $D_{\xi_{\text{st}}}$ may be assumed to look as in Figure 7, with two elliptic singular points along $\hat{K}$ (cf. [16, Prop. 4.8.11]).

Given two such knots K, K' , which coincide outside a compact set, we may assume the corresponding discs D, D' to coincide inside a small ball B^0 as indicated by a dashed circle in Figure 7. Notice that one may assume the characteristic foliation $(\partial B^0)_{\xi_{\text{st}}}$ to be the standard one with two singular elliptic points only and no closed orbits. Since the characteristic foliations $D_{\xi_{\text{st}}}$ and $D'_{\xi_{\text{st}}}$ determine the germ of the contact structure, we find balls B, B' containing B^0 and D, D' , respectively, and a contactomorphism $(B, \xi_{\text{st}}) \rightarrow (B', \xi_{\text{st}})$ fixed on B^0 and sending $\hat{K}$ to $\hat{K}'$ (and D to D'). By Eliashberg's classification of tight contact structures on the 3-ball [9, Thm. 2.1.3], this contactomorphism extends to a contactomorphism of (S^3, ξ_{st}) . Finally, by the parametric version of Eliashberg's result and the fact that the relative diffeomorphism group $\text{Diff}(D^3, S^2)$ is connected [1], this contactomorphism of (S^3, ξ_{st}) is contact isotopic to the identity relative to B^0 . This translates into a Legendrian isotopy between the long knots K and K' . $\square$

Remark. There is a slightly roundabout but more elementary argument for the final step in the preceding proof. We may choose a smaller ball B^1 inside B^0 sharing all its properties, and contained in the neighbourhood νL_0 of a standard Legendrian unknot L_0 inside B^0 . We then have to deal with a contactomorphism of (S^3, ξ_{st}) fixed on νL_0 . A result of Giroux [18] can then be used to show that such a contactomorphism is contact isotopic rel νL_0 (and in particular rel B^1) to the identity, see [5, p. 161/2].

5. CANCELLATION OF 1- AND 2-HANDLES

As described in [19, Section 2] or [20, Section 11.1], the diagram in Figure 8, depicting a 1-handle attached to D^4 along $S^3 = \partial D^4$, represents the manifold $S^1 \times S^2$ with its standard Stein fillable contact structure. Since contact (-1) -surgery along a Legendrian knot preserves Stein fillability, the diagram in Figure 9 represents a Stein fillable contact structure on S^3 , so this is a diagram for (S^3, ξ_{st}) . A slightly stronger formulation of this observation is that the 3-ball bounded by a 2-sphere S around this configuration (indicated by a dashed line) carries the tight contact structure uniquely determined, up to isotopy, by the characteristic foliation $S_{\xi_{\text{st}}}$.


FIGURE 8. $S^1 \times S^2$ with its standard tight contact structure.


FIGURE 9. Cancelling handle pair.

We now want to tie up this observation with Propositions 2 and 3 in order to prove the following result, which tells us how to replace 1-handles by contact $(+1)$ -surgeries along trivial Legendrian knots. This corresponds to the ‘dotted circle’ notation in Kirby diagrams [20, p. 167/8]; see also [27, p. 167/8] for a related discussion. (It is interesting to observe the analogy on the level of handlebodies: a dotted circle indicates a 2-handle being ‘dug out’ from D^4 , while a contact $(+1)$ -surgery corresponds to attaching a 2-handle to the concave end of a symplectic cobordism.)

The first diagram described in the next theorem is said to be in ‘standard form’; by [19, Thm. 2.2] any diagram involving 1-handles can be written in that way.

Theorem 4. *The two surgery diagrams in Figure 10 represent contactomorphic contact manifolds. Here it is understood that the surgery coefficients (relative to the contact framing) of the curves making up the Legendrian tangle remain unchanged.*

Proof. By the cancellation lemma [3, Prop. 8], cf. [16, Prop. 6.4.5], the first two diagrams shown in Figure 11 are equivalent. The Legendrian knots in the second diagram are Legendrian isotopic, relative to the ball indicated by a dashed ellipse, to the knots in the third diagram by Propositions 2 and 3. Finally, the fourth diagram in Figure 11 is obtained from the third by cancelling a 1-handle with a 2-handle as described above. The theorem follows by applying this procedure to each 1-handle. $\square$

FIGURE 10. Replacing 1-handles by $(+1)$ -surgeries.

When we are dealing with a link diagram where each strand passing over a 1-handle is stabilised, one can give a more direct proof of Theorem 4, using only handle slides and the cancellation lemma from [4], cf. [16, Prop. 6.4.5], but none of the awkward arguments involving relative isotopies. This is illustrated in Figure 12 for a single knot L ; the argument for several stabilised strands passing over a given 1-handle is analogous. (In this sequence of pictures, ‘ $\cong$ ’ indicates a contactomorphism; an arrow, a Legendrian isotopy.) First we introduce a pair of cancelling contact (± 1) -surgeries along the Legendrian knots $L_{\pm}$. Then apply Proposition 2 to turn L_+ into a meridian of L_- . Now form $L \#_{Lb} L_-$ as in Proposition 1. The obvious Legendrian isotopy that gives the fifth picture in the sequence of Figure 12 is a combination of what are called ‘move 4’ and ‘move 5’ in [19]; see Section 7 below. To get the final picture, apply the cancellation from Figure 9.

The particular positioning of the meridional $(+1)$ -surgery curve is irrelevant, see the following section. The requirement that L be stabilised is not restrictive; this will be discussed in Section 8.

6. SLIDING 1-HANDLES

An easy way to realise 1-handle slides in a Kirby diagram is to represent the 1-handle over which one wants to slide another 1-handle by a dotted circle, see [20, Figure 5.39]. In the Legendrian setting, at first sight there appears to be less flexibility in performing such moves, because it may not be possible to move one of the attaching balls of the sliding handle through the dotted circle by a Legendrian isotopy. However, one can in fact ensure complete flexibility by moving the dotted circle, which in the contact setting corresponds to a $(+1)$ -surgery along a Legendrian




FIGURE 11. Equivalent surgery diagrams.




FIGURE 12. Replacing 1-handles by (+1)-surgeries in stabilised case.

meridian. Figures 13 to 15 show Legendrian isotopies of such a meridian; Figure 16 illustrates that a 1-handle slide of either orientation can be realised by a contact isotopy.




FIGURE 13. Isotopic Legendrian meridians I.




FIGURE 14. Isotopic Legendrian meridians II.




FIGURE 15. Isotopic Legendrian meridians III.




FIGURE 16. 1-handle slides of either orientation.

7. MOVES 4 AND 5

In addition to the three usual Legendrian Reidemeister moves, in the presence of 1-handles there are three further moves to consider, see [19, Thm. 2.2]. We now describe these moves when the 1-handle has been replaced by a contact $(+1)$ -surgery along a Legendrian unknot. Figures 17 and 18 give the simple analogue of move 4 and 5, respectively; move 6 will be discussed in Section 8. Observe that moves 4 and 5 are Legendrian isotopies in the complement of the unknot, so the surgery plays no role here. The isotopy between the first and the second line in Figure 18 is given by two Legendrian Reidemeister moves of the second kind (involving only the two strands passing through the unknot) and a couple of Reidemeister moves of the third kind (involving in addition the unknot).




FIGURE 17. Move 4.




FIGURE 18. Move 5.

8. LIGHT BULB TRICK, MOVE 6, AND ROLFSEN TWIST

The requirement at the end of Section 5 that the knot passing over the 1-handle be stabilised turns out not to be any restriction at all. This is seen by a move that is essentially the light bulb trick [31, p. 257], see Figure 19. In the contact geometric setting, this move has been observed earlier by Gompf [19, Figure 18]. The move from the first to the second line in Figure 19 is ‘move 6’ in [19].

Remark. Write $S_{\pm}L$ for the positive or negative stabilisation, respectively, of an oriented Legendrian knot L , corresponding to adding a down or up zigzag to the front projection of L . Denote by

$$\xi_{\text{st}} = \ker(z d\theta + x dy - y dx).$$

the standard tight (and Stein fillable) contact structure on $S^1 \times S^2 \subset S^1 \times \mathbb{R}^3$. Then the light bulb trick says that any Legendrian knot L in $(S^1 \times S^2, \xi_{\text{st}})$ that crosses some sphere $\{\theta_0\} \times S^2$ exactly once is Legendrian isotopic to its double stabilisation S_+S_-L . Observe that ξ_{st} is trivial as a 2-plane bundle, with trivialisation given by the vector fields $x\partial_\theta - z\partial_y + y\partial_z$ and $y\partial_\theta + z\partial_x - x\partial_z$.

Contrast this with [33, Thm. 4.1.1], which says that with ξ_{st} replaced by any contact structure of non-zero Euler class, the knots L and S_+S_-L are not Legendrian isotopic, in spite of being Legendrian regularly homotopic and framed isotopic; they can be distinguished by a Vassiliev invariant.




FIGURE 19. The light bulb trick.

We now want to show how this move 6 (and hence the light bulb trick) can be performed when the 1-handle has been replaced by a $(+1)$ -surgery along a Legendrian unknot L_0 . Figure 20 shows that move 6 translates into a handle slide as in Proposition 1, where the connected sum $L \# L_0^+$ is not taken along a cusp, but along a pair of parallel strands; cf. [14, Figure 2] for this connected sum. A connected sum of this kind with a Legendrian unknot can be undone by a first Reidemeister move.




FIGURE 20. Move 6 via connected sum.

The following alternative approach to this move, as illustrated in Figure 21, may be worth noting. Here the construction is based on the fact that L_0^+ is a Legendrian unknot in the surgered manifold. Hence, we may assume that, in the surgered manifold, L_0^+ bounds a disc foliated by Legendrian curves as in Figure 7. This allows us to move the upper strand of L_0^+ to the lower one by a Legendrian isotopy across this new meridional disc in the surgered manifold, as shown in the second step in the sequence of Figure 21; the remaining isotopies are given by Reidemeister moves.




FIGURE 21. Move 6 via slide across meridional disc in surgered manifold.

Both versions of handle addition used here for performing move 6 should be regarded as incarnations of the second Kirby move.

At first sight, it is a little surprising that if L is a Legendrian knot passing over a 1-handle in a description of some contact manifold (M, ξ) , then there is a contactomorphism of (M, ξ) sending L to any of its stabilisations $S_+^k S_-^m L$. This can be seen from Figure 12 by replacing L_- with $S_+^k S_-^m L_-$, and L_+ with the Legendrian push-off of $S_+^k S_-^m L_-$.

When we replace the 1-handle in Figure 12 by a contact $(+1)$ -surgery along a Legendrian unknot L_0 , linked once with L , this contactomorphism may be regarded as a Rolfsen twist along L_0 (see [20, Figure 5.27] for the topological picture). Observe that the surgery coefficient of L_0 with respect to the surface framing of L_0 equals zero; this coefficient remains zero under a Rolfsen twist.

9. LONG LEGENDRIAN KNOTS

We now return to the completion procedure $K \mapsto \hat{K}$ for long Legendrian knots K , where we regard $\hat{K}$ as a Legendrian knot in (S^3, ξ_{st}) . The following theorem answers a question of Fuchs and Tabachnikov [15, Section 3.5].

Theorem 5. *The completion map $K \mapsto \hat{K}$ sets up a one-to-one correspondence between Legendrian isotopy classes of long Legendrian knots in $(\mathbb{R}^3, \xi_{\text{st}})$ and those of Legendrian knots in (S^3, ξ_{st}) .*

Proof. The map $K \mapsto \hat{K}$ induces a surjective map $[K] \mapsto [\hat{K}]$ on Legendrian isotopy classes. We need to show that this last map is injective.

Thus, let K_0 and K_1 be two long Legendrian knots in $(\mathbb{R}^3, \xi_{\text{st}})$ whose completions $\hat{K}_0, \hat{K}_1$ are Legendrian isotopic in (S^3, ξ_{st}) . This Legendrian isotopy extends to a contact isotopy of (S^3, ξ_{st}) [16, Thm. 2.6.2]. Let ϕ_1 be the time-1-map of that isotopy, which we may assume to be the identity on an interval $I \subset \hat{K}_0 \cap \hat{K}_1$ containing the point $\infty := S^3 \setminus \mathbb{R}^3$. Our aim is to find a contactomorphism ϕ'_1 of $(\mathbb{R}^3, \xi_{\text{st}})$ sending K_0 to K_1 , and with $\phi'_1 = \text{id}$ outside a compact set, say some large ball B . For then, as at the end of the proof of Proposition 3, we see that ϕ'_1 is in fact contact isotopic to the identity relative to $\mathbb{R}^3 \setminus B$. This translates into a Legendrian isotopy between the long knots K_0 and K_1 .

We provide two constructions of such a contactomorphism ϕ'_1 . The first one is based on the contact disc theorem; the second construction, which is inspired by the proof of Lemma 5.3 in [14], uses the classification of tight contact structures on solid tori. We include the second argument because the method may have some interest beyond the present application.

In new local coordinates (x, y, z) around $\infty = (0, 0, 0)$ we may think of I as the interval $(0, t, 0)$, $|t| \leq \varepsilon$, and ξ_{st} as being given by $dz + x dy = 0$. This will be the set-up used for both arguments.

(1) Write $\phi_1(x, y, z) = (\tilde{x}, \tilde{y}, \tilde{z})$ and consider the contact dilation

$$\delta_s(x, y, z) = (sx, sy, s^2z).$$

Let B_∞ be a small closed ball around $\infty = (0, 0, 0)$ with $I \cap B_\infty$ contained in the interior of I . As shown in the proof of the contact disc theorem [16, Thm. 2.6.7], with $A := \frac{\partial \tilde{z}}{\partial \tilde{z}}(0) \in \mathbb{R}^+$ and $C := \frac{\partial \tilde{y}}{\partial x}(0)$ we obtain a 1-parameter family of contact embeddings of B_∞ by

$$(x, y, z; s) \mapsto \begin{cases} \delta_s^{-1} \circ \phi_1 \circ \delta_s(x, y, z) & , s \in (0, 1], \\ (Ax, Cx + y, Az - \frac{1}{2}ACx^2) & , s = 0. \end{cases}$$


These contact embeddings of B_∞ fix I pointwise, and the embedding for $s = 0$ is contact isotopic to the identity via contact embeddings fixing I pointwise. The contact isotopy extension theorem [16, Thm. 2.6.12] then yields the desired contactomorphism ϕ'_1 .

(2) For $\delta > 0$ sufficiently small, the embedded discs $\phi_1(D_{y_0})$, $|y_0| \leq \varepsilon$, with


$$D_{y_0} := \{(x, y_0, z) : x^2 + z^2 \leq \delta^2\},$$

define a foliation transverse to ∂_{y_0} . So for each fixed pair (x_0, z_0) , the holonomy map $y_0 \mapsto y'_0$ sending y_0 to the y -coordinate of the intersection of $\phi_1(D_{y_0})$ with the y -line through $(x_0, 0, z_0)$ defines, near $y_0 = 0$, a diffeomorphism onto its image.

This allows us to apply the 1-dimensional disc theorem [21, Thm. 8.3.1] with two parameters (x_0, z_0) in order to isotope ϕ_1 to a diffeomorphism that sends each disc D_{y_0} (possibly for some smaller δ) into the plane $\mathbb{R} \times \{y_0\} \times \mathbb{R}$, fixing the origin. Next we apply the 2-dimensional disc theorem with one parameter y_0 in order to isotope this diffeomorphism further to a diffeomorphism ϕ_2 with $\phi_2(K_0) = K_1$, $\phi_2 = \text{id}$ near ∞ and $\phi_2 = \phi_1$ outside a slightly bigger neighbourhood. Since ϕ_1 was a contactomorphism, and the contact framing of I is given by ∂_x , we may in fact assume — by the neighbourhood theorem for Legendrian curves — that ϕ_2 is also a contactomorphism of a neighbourhood of $\hat{K}_0$ onto a neighbourhood of $\hat{K}_1$.

FIGURE 22. The diffeomorphism ϕ_2 near ∞ .

So we have a situation as illustrated in Figure 22, with ϕ_2 a contactomorphism outside a solid torus $S^1 \times D^2$. The boundary T^2 of that solid torus is convex with characteristic foliation as shown in Figure 23; this has slope -1 (the dividing curves are shown as dashed lines).

FIGURE 23. The characteristic foliation on T^2 .

Up to isotopy rel boundary, there is a unique tight contact structure on the solid torus with this characteristic foliation on the boundary [22, Prop. 4.3]. So we find a contactomorphism from $(\phi_2(S^1 \times D^2), T\phi_2(\xi_{st}))$ to $(\phi_2(S^1 \times D^2), \xi_{st})$. Now extend this contactomorphism by the identity outside $\phi_2(S^1 \times D^2)$ to a contactomorphism $\phi_3: (S^3, T\phi_2(\xi_{st})) \rightarrow (S^3, \xi_{st})$. Then $\phi'_1 := \phi_3 \circ \phi_2$ has the desired properties. $\square$

10. DESTABILISATION AND THE FIRST KIRBY MOVE

A Kirby move of the first kind consists in adding to (or deleting from) a given surgery diagram an unknotted circle, not linked with any component of the given link, with framing ± 1 . This does not change the topology of the 3-manifold described by the surgery diagram.

There is a not entirely satisfying analogue of this construction for contact surgery diagrams describing an overtwisted contact 3-manifold. Here the move consists in adding the link in Figure 24 to the given diagram. This corresponds, as explained

in [6], to forming the connected sum with S^3 equipped with an overtwisted contact structure that is homotopic, as a 2-plane field, to ξ_{st} ; such a connected sum does not change a given overtwisted contact manifold by Eliashberg's theorem [8], cf. [16, Section 4.7]. The dashed circle indicates the boundary of an overtwisted disc, cf. [6, Fig. 2]. (In the best of all possible worlds one should expect that it were sufficient to add but the 'shark' on the left, corresponding to a single topologically (-1) -framed unknot. This, alas, changes the Hopf d_3 -invariant of the contact structure.)




FIGURE 24. The 'first Kirby move'.

Now suppose we are given a stabilised Legendrian knot in the original diagram. When we perform the connected sum of such a knot with the mentioned boundary of an overtwisted disc, we obtain (after a couple of second Reidemeister moves) the situation on the top left of Figure 25. In the first row of that figure we perform a Kirby move of the second kind; in the second and third row, a move as in Figure 15. The remaining moves are the obvious Legendrian Reidemeister moves. Thus, this figure gives an explicit demonstration of the well-known result that the connected sum of a Legendrian knot with the boundary of an overtwisted disc (disjoint from the given knot) leads to a destabilisation, cf. [16, pp. 316–319].

Another way of phrasing this result is that any Legendrian knot with overtwisted complement (such 'loose' knots will be the topic of the next section) is Legendrian isotopic to any of its stabilisations, with the appropriate number of shark-like meridional curves added (as well as the curves on the right of Figure 24). This corresponds to [7, Lemma 4.7].




FIGURE 25. Destabilisation via Kirby moves.

We leave it to the reader to verify the following simple observation. Both positive and negative stabilisations can be effected by a shark with his 'mouth' on the left.

One can always perform the stabilisation along a strand oriented towards the right, say, by first performing a first Legendrian Reidemeister move to create an up or down cusp on the right. The sign of the stabilisation affects the positioning of the shark-like meridian as shown in Figure 26.




FIGURE 26. Stabilisations and respective position of shark.

11. LOOSE KNOTS

A Legendrian knot L in a contact 3-manifold (M, ξ) is called *loose* if $\xi|_{M \setminus L}$ is overtwisted. The following theorem is a generalisation of a result due to K. Dymara [7]. When we speak of a Legendrian knot $L \subset (M, \xi)$ as a *framed* knot, we always mean the framing defined by the contact structure.

Theorem 6. *Let L_0, L_1 be loose Legendrian knots in a (closed, connected) contact 3-manifold (M, ξ) , with an overtwisted disc $\Delta \subset M \setminus (L_0 \cup L_1)$. Assume that:*

- (i-a) L_0 and L_1 are isotopic as framed knots;
- (i-b) the topological knot type of L_0, L_1 contains infinitely many non-isotopic framed knots;
- (ii-a) L_0 and L_1 are Legendrian regularly homotopic as Legendrian immersions;
- (ii-b) the connected component of L_0, L_1 in the space of framed immersions $S^1 \rightarrow M$ contains infinitely many components of the space of Legendrian immersions $S^1 \rightarrow (M, \xi)$.

Then L_0 and L_1 are Legendrian isotopic.

Remark. Condition (i-b) being satisfied for all knot types in M is equivalent to there being *no* summands $S^1 \times S^2$ in the prime decomposition of M , see the remark on p. 488 of [23] and, for a proof, [2, Thm. 2.0.5]. One direction of this equivalence is simple. The existence of such a summand is equivalent to M containing a non-separating 2-sphere; the framing of a knot K intersecting such a 2-sphere transversely in a single point can be changed by the (topological) light-bulb trick by any even number of twists, resulting in only two different *framed* isotopy classes within the isotopy class of K .

Condition (ii-b) is equivalent to there being *no* class $\alpha \in H_2(M)$ on which the Euler class of ξ evaluates non-trivially and that can be represented by a mapping $S^1 \times S^1 \rightarrow M$ whose restriction to the second factor is a loop freely homotopic to L_0, L_1 , see [33, Prop. 3.1.4]. In particular, this condition is satisfied when ξ is trivial as a plane bundle (as was assumed in [7]).

Proof of Theorem 6. By [4] we can represent (M, ξ) by a surgery diagram $\mathbb{L} = \mathbb{L}^- \sqcup \mathbb{L}^+$ in (S^3, ξ_{st}) , i.e. a Legendrian link $\mathbb{L}$ such that contact (-1) -surgery along the components of $\mathbb{L}^-$ and contact $(+1)$ -surgery along the components of $\mathbb{L}^+$ yields

(M, ξ) . Thus, we may think of L_0, L_1 as Legendrian knots in (S^3, ξ_{st}) disjoint from $\mathbb{L}$.

By assumption, there is an isotopy from L_0 to L_1 (as framed knots) in M . This translates into a composition of isotopies in $S^3 \setminus \mathbb{L}$ and handle slides over the 2-handles attached along $\mathbb{L}$. The latter can be performed as band-sums.

According to [15, Thm. 4.4], any topological isotopy between two Legendrian knots in (S^3, ξ_{st}) can be converted into a Legendrian isotopy of suitable stabilisations of the two knots. The argument given there remains valid inside $S^3 \setminus \mathbb{L}$. Similarly, any topological band-sum can be performed as a Legendrian band-sum, provided we permit ourselves to stabilise one of the summands (in our case: L_0) as well as the Legendrian band, where stabilisations are necessary for realising left-handed twists in the band.

Thus, we can Legendrian isotope a suitable stabilisation $S_+^k S_-^m L_0$ to some stabilisation $S_+^l S_-^n L_1$. Since every stabilisation adds a negative twist to the contact framing, conditions (i-a) and (i-b) imply that $k + m = l + n$. Similarly, conditions (ii-a) and (ii-b) imply that $k - m = l - n$. This follows from the fact that the Legendrian regular homotopy class of a parametrised closed curve $\gamma: S^1 \rightarrow (M, \xi)$ is determined by the homotopy class of the bundle map $T\gamma \otimes \mathbb{C}: TS^1 \otimes \mathbb{C} \rightarrow \xi$, cf. [16, Section 6.3], and the effect of a positive or negative stabilisation on this bundle map which, being a local modification, can be understood in the same way as the effect of a stabilisation on the rotation number. We conclude that $k = l$ and $m = n$.

The Legendrian isotopy between $S_+^k S_-^m L_0$ and $S_+^k S_-^m L_1$ extends to a contact isotopy ϕ_t , $t \in [0, 1]$, of (M, ξ) . Since $\phi_t(L_0)$, $t \in [0, 1]$, defines a Legendrian isotopy from L_0 to $\phi_1(L_0)$, we may replace L_0 in the sequel of the argument by $\phi_1(L_0)$. (Observe that we may construct ϕ_t in such a way that it remains stationary on the overtwisted disc Δ , so the complement of L_1 and this new L_0 still contains Δ .)

Thus, we may assume that $S_+^k S_-^m L_0$ and $S_+^k S_-^m L_1$ actually coincide for some specific choice of these stabilisations. At any rate, we also find some segment where L_0 and L_1 coincide. Since there is an overtwisted disc in the complement of $L_0 \cup L_1$, we may perform first Kirby moves in $M \setminus (L_0 \cup L_1)$ without changing the situation (up to contactomorphism). Thus, by the discussion in Section 10, the knots L_0, L_1 are Legendrian isotopic in (M, ξ) to their stabilisations $S_+^k S_-^m L_0 = S_+^k S_-^m L_1$ with the appropriate number of shark-like meridians added; those meridians can be added in a neighbourhood of the segment where L_0 and L_1 coincide and thus may be taken to be identical for L_0 and L_1 . This proves that L_0 and L_1 are Legendrian isotopic. $\square$

Remark. Conditions (i-a) and (ii-a) in Theorem 6 are obviously necessary. The examples in Section 4 of [33] show that (i-b) and (ii-b) are likewise necessary.

With the help of the contact surgery presentation theorem from [4] one can also formulate a result about isotopies of Legendrian knots L_0, L_1 in an arbitrary contact 3-manifold (M, ξ) . Think of (M, ξ) as being represented by a surgery diagram $\mathbb{L} = \mathbb{L}^- \sqcup \mathbb{L}^+$ in (S^3, ξ_{st}) , and of L_0, L_1 as knots in $S^3 \setminus \mathbb{L}$. We visualise L_0, L_1 and the link $\mathbb{L}$ in the front projection. Then L_0 and L_1 are Legendrian isotopic in (M, ξ) if and only if the front of L_0 can be turned into the front of L_1 by Legendrian Reidemeister moves of the first three kinds (in the complement of $\mathbb{L}$) and by second Kirby moves. This follows in analogy with the topological case and from the corresponding result for $\mathbb{L} = \emptyset$ [32]. Any Legendrian isotopy that does

not pass through the belt sphere of one of the 2-handles corresponding to $\mathbb{L}$ can be made disjoint from those 2-handles and thus takes place in $S^3 \setminus \mathbb{L}$; a Legendrian isotopy with a single transverse passage through one of the belt spheres corresponds to a second Kirby move.

Acknowledgements. Some of this research was carried out during a stay of F. D. at the Mathematical Institute of the Universität zu Köln, supported by grant no. 10631060 of the National Natural Science Foundation of China. H. G. is partially supported by DFG grant GE 1245/1-2 within the framework of the Schwerpunktprogramm 1154 “Globale Differentialgeometrie”.

REFERENCES

- [1] J. CERF, *Sur les difféomorphismes de la sphère de dimension trois* ($\Gamma_4 = 0$), Lecture Notes in Math. **53**, Springer-Verlag, Berlin (1968).
- [2] V. CHERNOV (TCHERNOV), Framed knots in 3-manifolds and affine self-linking numbers, *J. Knot Theory Ramifications* **14** (2005), 791–818.
- [3] F. DING AND H. GEIGES, Symplectic fillability of tight contact structures on torus bundles, *Algebr. Geom. Topol.* **1** (2001), 153–172.
- [4] F. DING AND H. GEIGES, A Legendrian surgery presentation of contact 3-manifolds, *Math. Proc. Cambridge Philos. Soc.* **136** (2004), 583–598.
- [5] F. DING AND H. GEIGES, Legendrian knots and links classified by classical invariants, *Commun. Contemp. Math.* **9** (2007), 135–162.
- [6] F. DING, H. GEIGES AND A. I. STIPSICZ, Surgery diagrams for contact 3-manifolds, *Turkish J. Math.* **28** (2004), 41–74.
- [7] K. DYMARA, Legendrian knots in overtwisted contact structures, [arXiv:math.GT/0410122](https://arxiv.org/abs/math.GT/0410122).
- [8] YA. ELIASHBERG, Classification of overtwisted contact structures on 3-manifolds, *Invent. Math.* **98** (1989), 623–637.
- [9] YA. ELIASHBERG, Contact 3-manifolds twenty years since J. Martinet’s work, *Ann. Inst. Fourier (Grenoble)* **42** (1992), 165–192.
- [10] YA. ELIASHBERG AND M. FRASER, Classification of topologically trivial Legendrian knots, in: *Geometry, Topology and Dynamics* (Montréal, 1995), CRM Proc. Lecture Notes **15**, American Mathematical Society, Providence (1998), 17–51.
- [11] YA. ELIASHBERG AND M. FRASER, Topologically trivial Legendrian knots, [arXiv:0801.2553](https://arxiv.org/abs/0801.2553).
- [12] J. B. ETNYRE, Legendrian and transversal knots, in: *Handbook of Knot Theory*, Elsevier, Amsterdam (2005), 105–185.
- [13] J. B. ETNYRE AND K. HONDA, Knots and contact geometry I: Torus knots and the figure eight knot, *J. Symplectic Geom.* **1** (2001), 63–120.
- [14] J. B. ETNYRE AND K. HONDA, On connected sums and Legendrian knots, *Adv. Math.* **179** (2003), 59–74.
- [15] D. FUCHS AND S. TABACHNIKOV, Invariants of Legendrian and transverse knots in the standard contact space, *Topology* **36** (1997), 1025–1053.
- [16] H. GEIGES, *An Introduction to Contact Topology*, Cambridge Stud. Adv. Math. **109**, Cambridge University Press (2008).
- [17] P. GHIGGINI, P. LISCA, AND A. I. STIPSICZ, Tight contact structures on some small Seifert fibered 3-manifolds, *Amer. J. Math.* **129** (2007), 1403–1447.
- [18] E. GIROUX, Sur les transformations de contact au-dessus des surfaces, in *Essays on Geometry and Related Topics*, eds. É. Ghys et al., vol. 2, Monogr. Enseign. Math. **38** (Genève, 2001), 329–350.
- [19] R. E. GOMPF, Handlebody construction of Stein surfaces, *Ann. of Math. (2)* **148** (1998), 619–693.
- [20] R. E. GOMPF AND A. I. STIPSICZ, *4-Manifolds and Kirby Calculus*, Grad. Stud. Math. **20**, American Mathematical Society, Providence (1999).
- [21] M. W. HIRSCH, *Differential Topology*, Grad. Texts in Math. **33**, Springer-Verlag, Berlin (1976).
- [22] K. HONDA, On the classification of tight contact structures I, *Geom. Topol.* **4** (2000) 309–368; erratum: Factoring nonrotative $T^2 \times I$ layers, *Geom. Topol.* **5** (2001) 925–938.

- [23] J. HOSTE AND J. H. PRZYTICKI, Homotopy skein modules of orientable 3-manifolds, *Math. Proc. Cambridge Philos. Soc.* **108** (1990), 475–488.
- [24] Y. KANDA, On the Thurston-Bennequin invariant of Legendrian knots and nonexactness of Bennequin’s inequality, *Invent. Math.* **133** (1998), 227–242.
- [25] J. KOTUS, M. KRYCH AND Z. NITECKI, Global structural stability of flows on surfaces, *Mem. Amer. Math. Soc.* **37** (1982), no. 261.
- [26] P. LISCA AND A. I. STIPSICZ, Ozsváth-Szabó invariants and tight contact three-manifolds I, *Geom. Topol.* **8** (2004), 925–945.
- [27] B. ÖZBAĞCI AND A. I. STIPSICZ, *Surgery on Contact 3-Manifolds and Stein Surfaces*, Bolyai Soc. Math. Stud. **13**, Springer-Verlag, Berlin (2004).
- [28] M. M. PEIXOTO, Structural stability on two-dimensional manifolds, *Topology* **1** (1962), 101–120.
- [29] M. M. PEIXOTO, On an approximation theorem of Kupka and Smale, *J. Differential Equations* **3** (1967), 214–227.
- [30] O. PLAMENEVSKAYA, Transverse knots, branched double covers and Heegaard Floer contact invariants, *J. Symplectic Geom.* **4** (2006), 149–170.
- [31] D. ROLFSEN, *Knots and Links*, Mathematics Lecture Series **7**, Publish or Perish, Berkeley (1976).
- [32] J. ŚWIĄTKOWSKI, On the isotopy of Legendrian knots, *Ann. Global Anal. Geom.* **10** (1992), 195–207.
- [33] V. TCHERNOV, Vassiliev invariants of Legendrian, transverse, and framed knots in contact three-manifolds, *Topology* **42** (2003), 1–33.

DEPARTMENT OF MATHEMATICS, PEKING UNIVERSITY, BEIJING 100871, P. R. CHINA
E-mail address: dingfan@math.pku.edu.cn

MATHEMATISCHES INSTITUT, UNIVERSITÄT ZU KÖLN, WEYERTAL 86–90, 50931 KÖLN, GERMANY
E-mail address: geiges@math.uni-koeln.de